

Chemotherapy Protocol

BREAST CANCER

CYCLOPHOSPHAMIDE-EPIRUBICIN (75)- FLUOROURACIL

(FE₇₅C)

Regimen

- Breast Cancer – Cyclophosphamide-Epirubicin (75)-Fluorouracil (FE₇₅C)

Indication

- Neo-adjuvant / adjuvant therapy of breast cancer
- Metastatic breast cancer
- WHO Performance status 0, 1, 2

Toxicity

Drug	Adverse Effect
Cyclophosphamide	Dysuria, haemorrhagic cystitis, taste disturbances
Epirubicin	Cardio-toxicity, urinary discolouration (red)
Fluorouracil	Diarrhoea, stomatitis

The adverse effects listed are not exhaustive. Please refer to the relevant Summary of Product Characteristics for full details.

Monitoring

Regimen

- FBC, U&E's and LFT's prior to each cycle.
- Ensure adequate cardiac function before starting treatment. Baseline LVEF should be measured, particularly in patients with a history of cardiac problems or in the elderly.
- Patients with complete or partial dihydropyrimidine dehydrogenase (DPD) deficiency are at increased risk of severe and fatal toxicity during treatment with fluorouracil. All patients should be tested for DPD deficiency before initiation (cycle 1) to minimise the risk of these reactions

Dose Modifications

The dose modifications listed are for haematological, liver and renal function only. Dose adjustments may be necessary for other toxicities as well.

In principle all dose reductions due to adverse drug reactions should not be re-escalated in subsequent cycles without consultant approval. It is also a general rule

for chemotherapy that if a third dose reduction is necessary treatment should be stopped.

Please discuss all dose reductions / delays with the relevant consultant before prescribing if appropriate. The approach may be different depending on the clinical circumstances. The following is a general guide only.

Haematological

Prior to prescribing the following treatment criteria must be met on day 1 of treatment.

Criteria	Eligible Level
Neutrophils	equal to or more than $1 \times 10^9/L$
Platelets	equal to or more than $100 \times 10^9/L$

Consider blood transfusion if patient symptomatic of anaemia or has a haemoglobin of less than 8g/dL

In the adjuvant / neo-adjuvant setting, always check with the relevant consultant before delaying or reducing the dose in response to a toxicity.

If counts on day one are below these criteria for neutrophil and/or platelets then delay treatment for seven days or until recovery of counts. Treatment should only be started when these levels are reached. On subsequent cycles, if the counts are below these levels on day one then delay treatment until these levels are reached and / or consider reducing the dose to 80% of the original dose. If a second episode of neutropenia / thrombocytopenia occurs or the time to reach the eligible level is longer than seven days consider changing treatment. If patients experience febrile neutropenia or treatment delay due to neutrophil of less than $0.5 \times 10^9/L$ or platelets less than $50 \times 10^9/L$ for more than seven days then reduce the dose to 80% of the original dose. If neutropenia or thrombocytopenia recurs, the dosage should be either further reduce to 50% of the original dose or stop treatment. Growth factors may be considered as an alternative approach particularly in the neo-adjuvant and adjuvant setting.

Kidney Impairment

Drug	Creatinine Clearance (ml/min)	Dose (% of original dose)
Cyclophosphamide	more than 20	100
	10-20	75
	Less than 10	50
Epirubicin	Dose reduce in severe impairment only	
Fluorouracil	Consider dose reduction in severe renal impairment only	

Liver Impairment

Drug	Recommendation	
Cyclophosphamide	Dose reduction may not be necessary	
	Bilirubin (umol/L)	Dose (% of original)
Epirubicin	24-51	50
	51-85	25
	85 or greater	Contra-indicated
	If AST 2-4xULN give 50% dose, if the AST greater than 4xULN then give 25% dose	

Drug	Bilirubin µmol/L		AST/ALT units	Dose (%of original dose)
Fluorouracil	Less than 85		Less than 180	100%
	More than 85	or	More than 180	CI
	In moderate hepatic impairment reduce the initial dose by one third. In severe hepatic impairment reduce initial dose by one half. These doses may be increased if no toxicity occurs			

Other

Dose reductions or interruptions in therapy are not necessary for those toxicities that are considered unlikely to be serious or life threatening. For example, alopecia, altered taste or nail changes.

Epirubicin

Discontinue epirubicin if cardiac failure develops.

Regimen

21 day cycle for 6 cycles

Drug	Dose	Days	Administration
Cyclophosphamide	600mg/m ²	1	Intravenous bolus
Epirubicin	75mg/m ²	1	Intravenous bolus
Fluorouracil	600mg/m ²	1	Intravenous bolus

Dose Information

- Cyclophosphamide will be dose banded in accordance with the national dose bands (20mg/ml PM)
- Epirubicin will be dose banded in accordance with the national dose bands (2mg/ml PM)
- The maximum lifetime cumulative dose of epirubicin is 900mg/m².
- Fluorouracil will be dose banded in accordance with the national dose bands (25mg/ml PM)

Extravasation

- Cyclophosphamide – neutral
- Epirubicin – vesicant
- Fluorouracil - inflammitant

Additional Therapy

- Antiemetics

15-30 minutes prior to chemotherapy;

- dexamethasone 8mg oral or equivalent intravenous dose
- ondansetron 8mg oral or intravenous

As take home medication

- dexamethasone 4mg twice a day for 3 days oral
- metoclopramide 10mg three times a day when required oral
- ondansetron 8mg twice a day for three days oral

- Mouthwashes according to local or national policy on the treatment of mucositis
- Gastric protection with a proton pump inhibitor or a H₂ antagonist may be considered in patients considered at high risk of GI ulceration or bleed.

References

- 1.Coskan U, Gunel N, Onuk E et al. Effect of different neoadjuvant chemotherapy regimens on locally advanced breast cancer. Neoplasma 2003; 50 (3): 210-216.
- 2.Levine MN, Bramwell VH, Pritchard KI et al. Randomised trial of intensive Cyclophosphamide, Epirubicin and fluorouracil chemotherapy compared with Cyclophosphamide, Methotrexate and fluorouracil in premenopausal women with node-positive breast cancer. J Clin Oncol 1998; 16(8):2651-2658.
- 3.Brufman G, Colajori E, Ghilezan N et al. Doubling Epirubicin dose intensity (100mg/m² versus 50mg/m²) in the FEC regimen significantly increases response rates. An international randomised phase III study in metastatic breast cancer. Anns of Oncol 1997; 8: 155-162.

REGIMEN SUMMARY

Cyclophosphamide-Epirubicin (75)-Fluorouracil (FE₇₅C)

Day One

1. Dexamethasone 8mg oral or equivalent intravenous dose
2. Ondansetron 8mg oral or intravenous
3. Epirubicin 75mg/m² intravenous bolus over 10 minutes
4. Fluorouracil 600mg/m² intravenous bolus over 10 minutes
5. Cyclophosphamide 600mg/m² intravenous bolus over 10 minutes

Take Home Medicines

6. Dexamethasone 4mg twice a day for 3 days oral starting on day two of the cycle
7. Metoclopramide 10mg three times a day when required oral
8. Ondansetron 8mg twice a day for three days oral starting on the evening of day one of treatment

DOCUMENT CONTROL

Version	Date	Amendment	Written By	Approved By
1.2	Nov 2020	Updated monitoring with DPD testing Dose banding updated Coding removed	Donna Kimber Pharmacy Technician	Rebecca Wills Pharmacist
1.1	November 2014	Header changed Toxicities removed Adverse effects tabulated ≥ removed and written in full Dose modification tabulated Hepatic impairment updated Regimen tabulated Metoclopramide dose changed to 10mg Bolus removed from intravenous bolus throughout text Mucositis recommendation changed OPCS codes updated Dexamethasone TTO clarified Ondansetron TTO clarified Disclaimer added	Donna Kimber Pharmacy Technician	Dr Debbie Wright Pharmacist
1	Aug 2011	None	Anna Bunch Pharmacist Dr Debbie Wright Pharmacist	Dr Ellen Copson Consultant Medical Oncologist Dr Caroline Archer Consultant Medical Oncologist

This chemotherapy protocol has been developed as part of the chemotherapy electronic prescribing project. This was and remains a collaborative project that originated from the former CSCCN. These documents have been approved on behalf of the following Trusts;

Hampshire Hospitals NHS Foundation Trust
NHS Isle of Wight
Portsmouth Hospitals NHS Trust
Salisbury Hospital NHS Foundation Trust
University Hospital Southampton NHS Foundation Trust
Western Sussex Hospitals NHS Foundation Trust

All actions have been taken to ensure these protocols are correct. However, no responsibility can be taken for errors which occur as a result of following these guidelines.