

Patient information factsheet

Recovering well after breast surgery (glue)

We have given you this factsheet because you have had breast surgery (using glue for wound closure). It explains what to expect after your operation and how you can help your body to recover well. We hope it will help to answer some of the questions you may have. If you have any other questions or concerns, please do ask. We will be happy to help.

Common symptoms after breast surgery

Pain and discomfort

Most women experience some mild pain and/or a tingling sensation in their breast for a few days after the operation. You can take over-the-counter pain relief, such as paracetamol or ibuprofen-based products, to help ease your symptoms. Always read the label or instructions before taking medication.

You can take your next dose of pain relief at _____ (time).

Bruising and swelling

Bruising is common in the first few weeks after your operation, but it will disappear over time. However, some women may then develop a hard, tense bruise around their wound (this is called a haematoma).

Swelling is also common in the first few weeks after your operation and will usually settle over time. However, some women may then develop a collection of fluid under their skin near their wound (this is called a seroma).

If either a haematoma or a seroma develops, please contact our breast care nurses or your GP for advice.

When to seek medical advice

Most women recover well after breast surgery. However, as with any operation, there can be complications.

Contact Bramshaw Women's Unit on **023 8120 6035** (up to two weeks after your operation) or contact your GP or NHS **111** if you experience:

- increasing breast pain and/or swelling
- a temperature (fever)
- a loss of appetite and vomiting (being sick)
- any difficulty passing urine or emptying your bladder, a burning or stinging sensation when passing urine or passing urine more often than usual
- a painful, red, swollen, hot leg or difficulty bearing weight on your legs (this may be caused by a deep vein thrombosis (DVT))
- difficulty breathing

Patient information factsheet

Recovering from your operation

Wound closure

Your wound has been secured using special medical glue.

A surgical drain (a small plastic tube attached to a small bag) may have been inserted into your wound to help drain away fluid that might gather up and cause an infection after your operation. If a surgical drain has been inserted, we will give you a separate information factsheet about this.

If you have any concerns with how your wound is healing, contact our breast care nurses during office hours (please see below) or Bramshaw Women's Unit out of hours.

Eating and drinking

You can eat and drink as normal after your operation.

Washing

You may bathe and shower as normal after your operation. However, you should not soak or scrub your wound with a flannel or a sponge. You should also gently pat dry the area around your wound with a clean towel.

If you had a general anaesthetic, you may feel dizzy for a couple of days after your operation so please have an adult around to help you, if needed, and keep the bathroom door unlocked.

Wearing a bra

You should wear a well-fitting, supportive bra after your operation. You may find it more comfortable to wear a bra all the time (day and night) while you recover.

Advice after an anaesthetic and/or sedation

For the first 48 hours after your operation, you may feel more sleepy than usual and your judgement may be impaired. During this time, you must not:

- drive (your insurance company will have its own conditions for when you are insured to start driving again, so it is important you check your policy)
- drink alcohol
- work
- exercise
- handle machinery (including an oven or hob)
- sign any legal documents

Contact us

If you have any questions or concerns, please contact us.

Bramshaw Women's Unit
Telephone: **023 8120 6035**

Breast care nurses
Telephone: **023 8120 3753** (Monday to Thursday, 8am to 4pm, and Friday, 8am to 1pm)

Useful links

Royal College of Surgeons of England
Website: **www.rcseng.ac.uk**

For a translation of this document, or a version in another format such as easy read, large print, Braille or audio, please telephone **0800 484 0135** or email **patientsupporthub@uhs.nhs.uk**

For help preparing for your visit, arranging an interpreter or accessing the hospital, please visit **www.uhs.nhs.uk/additionalsupport**